

Dynamics of Self-Gravity Wakes in Dense Planetary Rings I. Pitch Angle

Shugo Michikoshi¹, Akihiko Fujii², Eiichiro Kokubo¹, and Heikki Salo³

shugo.michikoshi@nao.ac.jp, akihiko.fujii2@ucalgary.ca, kokubo@th.nao.ac.jp, and
heikki.salo@oulu.fi

ABSTRACT

We investigate the dynamics of self-gravity wakes in dense planetary rings. In particular, we examine how the pitch angle of self-gravity wakes depend on ring parameters using N -body simulations. We calculate the pitch angles using the two-dimensional autocorrelation function of the ring surface density. We obtain the pitch angles for the inner and outer parts of the autocorrelation function separately. We confirm that the pitch angles are 15 to 30 degrees for reasonable ring parameters, which are consistent with previous studies. We find that the inner pitch angle increases with the Saturn-centric distance, while it barely depends on the optical depth and the restitution coefficient of ring particles. The increase of the inner pitch angle with the Saturn-centric distance is consistent with the observations of the A ring. The outer pitch angle does not have the clear dependence on any ring parameters and is about 10–15 degrees. This value is consistent with the pitch angle of spiral arms in collisionless systems.

Subject headings: planets: rings, methods: N -body simulations

1. Introduction

The main rings of Saturn are composed of numerous icy particles and exhibit many different patterns, including non-axisymmetric structures on sub-kilometer scales. Gravitational forces between particles form gravitationally bound clumps, while differential rotation tears them apart. Due to these competing processes, spatial structures called gravitational or self-gravity wakes appear (Colombo et al. 1976; Salo 1992a). Salo (1992a) performed the first investigations of self-gravity wakes using N -body simulations. The typical radial scale of the wakes is comparable to the critical wavelength of gravitational instability (Toomre 1964), which is around several tens to hundred meters. Their azimuthal scale is much longer than their radial scale.

¹ Division of Theoretical Astronomy, National Astronomical Observatory of Japan, Osawa, Mitaka, Tokyo 181-8588, Japan

² University of Calgary, Calgary, Alberta, Canada T2N 1N4

³ Department of Physics, Astronomy Division, University of Oulu, PO Box 3000, FI-90014, Finland

Even though individual wakes are tiny structures, together they can have detectable influence on the large-scale photometric properties of the rings. The A and B rings exhibit a remarkable asymmetric brightness variation that was first discovered by Camichel (1958), confirmed later by the ground-based observations (Ferrin 1975; Reitsema et al. 1976), and by the Voyager image data (Franklin et al. 1987; Dones et al. 1993). Salo & Karjalainen (2003) and Salo et al. (2004) carried out the detailed photometric modeling of self-gravity wakes with N -body simulation. They found that the self-gravity wakes in N -body simulation can account for the brightness dependence on the longitude and elevation in the Voyager and ground-based observations. This conclusion was confirmed by the Hubble Space Telescope and Cassini observations (French et al. 2007; Porco et al. 2008).

Though there have been many numerical N -body simulations of self-gravity wakes (Salo 1992b; Richardson 1994; Daisaka & Ida 1999; Daisaka et al. 2001; Robbins et al. 2010; Yasui et al. 2012), the theoretical framework for dense planetary rings with self-gravity wakes has not been explored in detail yet. It is often argued that wakes in dense rings are dynamically similar to the spiral structure in differentially rotating galactic disks (Colombo et al. 1976; Franklin et al. 1987). Julian & Toomre (1966) studied theoretically the spiral structure in disk galaxies. They found that when disk surface density is perturbed, non-axisymmetric density enhancements appear. This mechanism is called swing amplification (Toomre 1981). Based on the swing amplification, we can estimate the pitch angle of a spiral arm (Julian & Toomre 1966; Toomre 1981). The pitch angle is the angle between the tangents to a spiral arm and a perfect circle, which measures how tightly the spiral arm is wound. The swing amplification model predicts that the pitch angle for Keplerian rotation is about $10\text{--}15^\circ$, which is roughly consistent with that of the wakes in rings. Thus, it has been considered that the swing amplification plays an important role in the self-gravity wake formation in rings.

According to the swing amplification model, the pitch angle of the self-gravity wakes depends on the shear rate (Julian & Toomre 1966; Michikoshi & Kokubo 2014). However, the dependence of the pitch angle on other ring parameters is still not well understood. Observationally Hedman et al. (2007) investigated the dependence of the pitch angle on the Saturn-centric distance. They found that the pitch angle varies systematically with the Saturn-centric distance. In the A ring, the pitch angle is $18^\circ\text{--}24^\circ$, which is consistent with the previous numerical simulations (Salo et al. 2004). In the middle A ring, the pitch angle is almost constant 18° , but in the outer A ring, it increases with the Saturn-centric distances to 24° . In addition, Hedman et al. (2007) pointed out that the pitch angle has the local maximum at the location of the strong density waves. From the Cassini UVIS occultation studies, Colwell et al. (2007) obtained the consistent results that the pitch angle in the A ring is $10\text{--}50^\circ$ and it increases with the Saturn-centric distance. They also found that the pitch angle in the B ring decreases with the Saturn-centric distance. From the HST observations, French et al. (2007) showed the increase of the pitch angle with the Saturn-centric distance in the A ring, which is consistent with the results in Colwell et al. (2007). The pitch angle dependence on the Saturn-centric distance has a complex structure. It is not a monotonic

function of the Saturnicentric distance. However, roughly speaking, the pitch angle decreases with the Saturnicentric distance in the B ring, while it increases in the A ring.

Salo (1995) performed N -body simulations and found that the wake pitch angle is about 20° from the autocorrelation function. Salo et al. (2004) found that the pitch angle varies with the autocorrelation distance. The pitch angle for small autocorrelation distance is about $25\text{--}30^\circ$, and decreases with the distance towards $15\text{--}20^\circ$. Using the photometric model they found that the longitude of the minimum brightness with respect to ansa increases from 17° to 23° with the elevation angle of the observer with respect to the ring plane. They argued that the observed minimum longitude for high elevation angles corresponds to the pitch angle for small distance, and that for low elevation angles corresponds to the pitch angle for large distance since the brightness for high elevation angles is dominated by the inner dense part of the wakes. However, the systematic study of the dependence of the pitch angle on ring parameters has not been conducted yet. Therefore, we perform such a survey with high resolution N -body resolutions. We calculate the pitch angle from the autocorrelation function of the ring surface density and investigate its dependence on the restitution coefficient of ring particles, the dynamical optical depth, and the Hill radius of a particle pair relative to the sum of their physical radii.

The outline of this paper is as follows. We describe the simulation model in section 2 and explain the calculation method of the pitch angle in section 3. In section 4, we present the results of the dependence of the pitch angle on ring parameters. Sections 5 and 6 are devoted for discussion and a summary, respectively.

2. Simulation Method

2.1. Equation of Motion

We adopt the shearing box method developed for local ring dynamics (e.g., Wisdom & Tremaine 1988; Michikoshi & Kokubo 2011). We introduce the rotating local Cartesian coordinates (x, y, z) . The x -axis points radially outward, the y -axis points in the direction of orbital motion, and the z -axis points to the normal to the ring plane. We assume $m_i \ll M_S$, $|x_i|, |y_i|, |z_i| \ll a$, and $|dx_i/dt|, |dy_i/dt|, |dz_i/dt| \ll a\Omega$, where M_S is the Saturn mass, m_i is the mass of i th ring particle, a is the Saturnicentric distance, and Ω is the orbital frequency. Then the equation of motion for the i th particle is (e.g., Petit & Henon 1986; Nakazawa & Ida 1988)

$$\frac{d^2x_i}{dt^2} = 2\Omega\frac{dy_i}{dt} + 3\Omega^2x_i + \sum_{j \neq i} \frac{Gm_j}{r_{ij}^3}(x_j - x_i), \quad (1)$$

$$\frac{d^2y_i}{dt^2} = -2\Omega\frac{dx_i}{dt} + \sum_{j \neq i} \frac{Gm_j}{r_{ij}^3}(y_j - y_i), \quad (2)$$

$$\frac{d^2 z_i}{dt^2} = -\Omega^2 z_i + \sum_{j \neq i} \frac{G m_j}{r_{ij}^3} (z_j - z_i), \quad (3)$$

where r_{ij} is the distance between i th and j th particles. The first terms in Equations (1) and (2) are the Coriolis force. The second term in Equation (1) is the tidal force, the first term in Equation (3) is the vertical force due to Saturn’s gravity, and the last terms in Equations (1)–(3) are the mutual gravity between ring particles.

The simulation box has a square shape with length L on the x - y plane. We use the sliding cell method where when a particle crosses a box boundary outward, the corresponding particle immediately comes into the box from the opposite boundary (Wisdom & Tremaine 1988; Toomre & Kalnajs 1991). The length L should be sufficiently larger than the characteristic scale length, the critical wavelength of the gravitational instability

$$\lambda_{\text{cr}} = \frac{4\pi^2 G \Sigma_0}{\Omega^2}, \quad (4)$$

where Σ_0 is the mean surface density (Toomre 1964). We adopt $L = 10\lambda_{\text{cr}}$.

For the collision model we adopt the constant restitution coefficient ϵ that ranges from 0.1 to 0.6. For $\epsilon > 0.7$, the velocity dispersion of ring particles increases monotonically and self-gravity wakes do not appear (Goldreich & Tremaine 1978). The effect of the velocity-dependent restitution coefficient is discussed in Section 4.1. For the velocity-dependent restitution coefficient model, we use the standard model of Bridges et al. (1984):

$$\epsilon(v_n) = \min(1, (v_n/v_c)^{-0.234}), \quad (5)$$

where v_n is the normal component of the relative velocity in collisions and $v_c = 0.0077 \text{ cm s}^{-1}$. We adopt the hard-sphere model as the collision model (e.g., Wisdom & Tremaine 1988; Salo 1991; Richardson 1994). In the hard-sphere model, when collision between particles is detected, the particle velocities change instantly. In this study since we neglect the friction between particles, the relative tangential velocity is conserved in collision and the effects of spin are not considered.

The equation of motion is integrated with a leapfrog integrator with timestep $\Delta t = (2\pi/\Omega)/200$. We calculate the self-gravity by direct summation using the special-purpose computer, GRAPE-DR (Makino et al. 2007). We introduce the cutoff of self-gravity by the subregion method (Daisaka & Ida 1999). In this method, the cutoff length corresponds to the box size L .

2.2. Ring Model

Ring particles are assumed to be identical for the sake of simplicity except for Section 4.4. For the identical particle model, all particles have the same mass $m_i = m_p$. This assumption is unrealistic since many observations show a power-law size distribution of ring particles (Zebker et al. 1985; French & Nicholson 2000). The effect of the realistic size distribution is discussed in

Section 4.4 where we adopt the power-law size distribution (Zebker et al. 1985; French & Nicholson 2000)

$$\frac{dN}{dR_p} \propto R_p^{-q}, \quad (6)$$

where R_p is the particle radius and q is the power-law index. From the Voyager radio occultation measurements, the power-law index increases with the Saturncentric distance from 2.7 to 3.0 in the A ring (Zebker et al. 1985). The stellar occultation observations showed the consistent results that the power-law indexes are $q = 2.75$ and 2.9 in the inner and outer A rings, respectively (French & Nicholson 2000). Considering these results, we choose the intermediate value $q = 2.8$. We set the lower and upper size limits as $R_{\min} = 150/\sqrt{\gamma}$ cm and $R_{\max} = 150\sqrt{\gamma}$ cm where $\gamma = R_{\max}/R_{\min} = 1$ –10.

For a ring with identical particles, the dynamical optical depth is given by

$$\tau = \frac{N\pi R_p^2}{L^2} = \frac{3\Sigma_0}{4R_p\rho_p} = 0.417 \left(\frac{\Sigma_0}{50 \text{ g cm}^{-2}} \right) \left(\frac{R_p}{100 \text{ cm}} \right)^{-1} \left(\frac{\rho_p}{0.9 \text{ g cm}^{-3}} \right)^{-1}, \quad (7)$$

where N is the number of particles in the simulation box and ρ_p is the particle density. For a low optical depth ring, the dynamical optical depth is equal to the observed optical depth (Salo & Karjalainen 2003; Robbins et al. 2010), while for a high optical depth ring, it is larger than the observed optical depth. Though the particle density of solid ice is 0.9 g cm^{-3} , ρ_p is estimated to be 0.45 g cm^{-3} from the best fit to the observations of azimuthal brightness variation in the A ring (Salo et al. 2004; French et al. 2007; Porco et al. 2008). The surface density is inferred from density waves, which is typically $\Sigma_0 \simeq 45 \text{ g cm}^{-2}$ in the A ring (e.g. Tiscareno et al. 2007). The surface density, radius, and particle density of ring particles collectively affect the structure of self-gravity wakes. We vary the dynamical optical depth from 0.3 to 1.8 in the identical particle models, which corresponds to the surface density of $\Sigma_0 = 36$ –216 g cm^{-2} with $R_p = 100$ cm and $\rho_p = 0.9 \text{ g cm}^{-3}$.

Both the semi-major axis and particle density affect the spatial structure. If the distance of the simulation box from the planet, i.e., the semi-major axis is large, the tidal force becomes less effective, allowing ring particles to gravitationally accumulate. In the region outside the Roche limit, particles can form gravitationally-bound aggregates. The particle density affects the strength of the self-gravity. Larger density makes particles pull each other more strongly, causing more distinct wakes. These factors are combined into one non-dimensional parameter, the Hill radius of a particle pair relative to the sum of their physical radii, defined as

$$\tilde{r}_H \equiv \frac{R_H}{2R_p} = 0.82 \left(\frac{M_S}{5.69 \times 10^{29} \text{ g}} \right)^{-1/3} \left(\frac{\rho_p}{0.9 \text{ g cm}^{-3}} \right)^{1/3} \left(\frac{a}{10^5 \text{ km}} \right), \quad (8)$$

where R_H is the Hill radius of the particle $R_H = (2m_p/3M_S)^{1/3}a$. With the particle density ρ_p fixed, $\tilde{r}_H$ is proportional to a . We vary the ratio from 0.68 to 1.06.

We introduce a parameter, initial Toomre Q to determine the initial random velocity of particles,

$$Q_{\text{ini}} = \frac{\sigma_x \Omega}{3.36 G \Sigma_0}, \quad (9)$$

where σ_x is the initial radial velocity dispersion (Toomre 1964). Since the ring relaxes into the stationary state quickly, Q_{ini} does not affect the final state (Salo 2012). Thus, we adopt $Q_{\text{ini}} = 2$.

The ring models are summarized in Table 1 for identical particles and in Table 2 for particles with size distributions. Note that the dynamical properties of rings with the identical particles are characterized by the three non-dimensional quantities: τ , $\tilde{\tau}_{\text{H}}$ and ϵ (e.g., Karjalainen & Salo 2004). The dimensional quantities such as particle size are irrelevant to simulation results. However, in order to compare with the actual rings, we add the corresponding dimensional quantities to Table 1 assuming $R_{\text{p}} = 100 \text{ cm}$ and $\rho_{\text{p}} = 0.45\text{--}0.90 \text{ g cm}^{-3}$. The surface density Σ_0 is calculated by Equation (7) and the Saturnicentric distance a is calculated by Equation (8).

Table 1. Ring Parameters and Pitch Angles for Identical Particle Models

Model	τ	$\tilde{r}_H$	ϵ	N	Σ_0 (g cm^{-2})	a (10^5 km)	L (km)	θ_i (deg.)	θ_o (deg.)	x_b (λ_{cr})	$ y_b $ (λ_{cr})
1	0.60	0.81	0.50	44156	36–72	0.78–0.99	0.48	25.8 ± 4.1	13.3 ± 4.2	0.26	0.53
2	0.70	0.81	0.10	70119	42–84	0.78–0.99	0.56	27.3 ± 6.0	10.9 ± 6.2	0.28	0.54
3	0.70	0.81	0.20	70119	42–84	0.78–0.99	0.56	25.7 ± 4.3	15.4 ± 7.0	0.27	0.55
4	0.70	0.81	0.30	70119	42–84	0.78–0.99	0.56	26.0 ± 4.3	13.8 ± 5.4	0.31	0.63
5	0.70	0.81	0.40	70119	42–84	0.78–0.99	0.56	25.7 ± 3.9	15.6 ± 4.7	0.23	0.49
6	0.70	0.81	0.50	70119	42–84	0.78–0.99	0.56	25.2 ± 3.1	15.1 ± 5.3	0.26	0.55
7	0.70	0.81	0.60	70119	42–84	0.78–0.99	0.56	27.0 ± 3.1	14.1 ± 3.3	0.14	0.28
8	0.70	0.81	$\epsilon(v_n)$	70119	42–84	0.78–0.99	0.56	24.5 ± 4.2	13.9 ± 4.1	0.27	0.60
9	0.80	0.60	0.50	17290	48–96	0.58–0.73	0.26	17.9 ± 1.5	10.0 ± 1.5	0.74	2.29
10	1.00	0.60	0.50	33770	60–120	0.58–0.73	0.33	22.1 ± 2.3	13.5 ± 1.5	0.04	0.10
11	1.20	0.60	0.50	58355	72–144	0.58–0.73	0.39	9.7 ± 2.1	17.5 ± 6.1	0.24	1.39
12	1.40	0.60	0.50	92666	84–168	0.58–0.73	0.46	8.2 ± 3.7	19.8 ± 10.5	0.19	1.28
13	1.60	0.60	0.50	138324	96–192	0.58–0.73	0.52	0.5 ± 2.7	-0.3 ± 2.3	0.03	3.98
14	1.80	0.60	0.50	196950	108–216	0.58–0.73	0.59	9.8 ± 1.4	17.8 ± 6.5	0.29	1.70
15	0.30	0.81	0.50	5519	18–36	0.78–0.99	0.24	30.2 ± 3.0	13.5 ± 2.6	0.29	0.50
16	0.40	0.81	0.50	13083	24–48	0.78–0.99	0.32	30.3 ± 2.5	13.2 ± 3.4	0.22	0.37
17	0.50	0.81	0.50	25553	30–60	0.78–0.99	0.40	29.5 ± 2.8	12.3 ± 4.4	0.21	0.37
18	0.70	0.81	0.50	70119	42–84	0.78–0.99	0.56	25.2 ± 3.1	15.1 ± 5.3	0.26	0.55
19	0.80	0.81	0.50	104667	48–96	0.78–0.99	0.64	25.9 ± 3.9	16.9 ± 6.0	0.16	0.32
20	0.90	0.81	0.50	149028	54–108	0.78–0.99	0.72	24.5 ± 3.9	12.8 ± 5.8	0.35	0.76
21	1.00	0.81	0.50	204428	60–120	0.78–0.99	0.80	24.5 ± 4.8	13.4 ± 5.7	0.30	0.67
22	1.10	0.81	0.50	272094	66–132	0.78–0.99	0.88	23.4 ± 4.1	13.6 ± 6.3	0.47	1.09
23	1.20	0.81	0.50	353253	72–144	0.78–0.99	0.96	23.7 ± 3.9	11.2 ± 6.2	0.54	1.23
24	1.30	0.81	0.50	449130	78–156	0.78–0.99	1.04	24.7 ± 4.8	13.6 ± 7.3	0.37	0.81
25	0.30	0.68	0.50	1932	18–36	0.66–0.83	0.14	26.9 ± 5.2	8.4 ± 5.2	0.80	1.58
26	0.30	0.73	0.50	2957	18–36	0.71–0.89	0.18	26.4 ± 3.6	13.5 ± 2.6	0.44	0.89
27	0.30	0.77	0.50	4073	18–36	0.75–0.94	0.21	28.1 ± 2.9	14.0 ± 3.3	0.30	0.57
28	0.30	0.81	0.50	5519	18–36	0.78–0.99	0.24	30.2 ± 3.0	13.5 ± 2.6	0.29	0.50
29	0.30	0.85	0.50	7370	18–36	0.82–1.04	0.28	32.2 ± 2.7	12.5 ± 3.2	0.30	0.47
30	0.30	0.90	0.50	10386	18–36	0.87–1.10	0.33	37.1 ± 3.8	13.1 ± 4.0	0.27	0.36
31	0.30	0.94	0.50	13482	18–36	0.91–1.15	0.38	42.3 ± 4.0	13.8 ± 4.1	0.24	0.26
32	0.30	0.98	0.50	17312	18–36	0.95–1.20	0.43	47.3 ± 5.3	13.3 ± 5.0	0.24	0.22
33	0.30	1.02	0.50	22008	18–36	0.99–1.24	0.48	50.4 ± 6.0	10.4 ± 5.2	0.31	0.25
34	0.30	1.06	0.50	27722	18–36	1.03–1.29	0.54	53.7 ± 5.5	7.3 ± 4.8	0.41	0.30
35	0.60	0.68	0.50	15457	36–72	0.66–0.83	0.28	22.1 ± 1.3	12.5 ± 1.6	0.20	0.50
36	0.60	0.73	0.50	23660	36–72	0.71–0.89	0.35	26.4 ± 1.9	13.3 ± 2.4	0.11	0.22
37	0.60	0.77	0.50	32585	36–72	0.75–0.94	0.41	25.2 ± 2.3	13.4 ± 4.0	0.18	0.39
38	0.60	0.85	0.50	58965	36–72	0.82–1.04	0.56	27.9 ± 4.3	13.7 ± 6.4	0.32	0.61
39	0.60	0.90	0.50	83088	36–72	0.87–1.10	0.66	32.0 ± 5.6	9.8 ± 6.3	0.40	0.64
40	0.60	0.94	0.50	107858	36–72	0.91–1.15	0.75	35.7 ± 5.2	16.7 ± 6.0	0.25	0.35

3. Structure of Self-Gravity Wakes

3.1. Formation of Self-Gravity Wakes

Figure 1 presents an example of the self-gravity wake formation for the standard model (model 1). The parameters are $\tau = 0.6$, $\tilde{r}_H = 0.81$, and $\epsilon = 0.5$. Initially particles are distributed uniformly in the simulation box (Fig.1a).

Figure 2 shows the time evolution of Q . The Q value is calculated from Equation (9), where we calculate σ_x from the root mean square of the radial velocity dx/dt . Due to inelastic collisions between particles the kinetic energy decreases and the velocity dispersion and Q decrease. When Q becomes $Q \simeq 1$ –2, the non-axisymmetric perturbations grow significantly (Julian & Toomre 1966; Toomre 1981). Accordingly, the condition for the emergence of the self-gravity wakes is obtained from $Q \lesssim 2$ (Salo 1995; Ohtsuki & Emori 2000; Daisaka et al. 2001)

$$\tau \gtrsim \begin{cases} 0.08\tilde{r}_H^{-3} & \text{for } \tilde{r}_H \lesssim 0.5 \\ 0.2\tilde{r}_H^{-3/2} & \text{for } \tilde{r}_H \gtrsim 0.5 \end{cases} . \quad (10)$$

When this condition is satisfied, the self-gravity wakes appear finally regardless of Q_{ini} (Salo 2012). As shown in Fig.1b, self-gravity wakes appear due to gravitational instability at $\tilde{t} = 1.0$ where $\tilde{t} = t/(2\pi/\Omega)$. Then the radial spacing of the self-gravity wakes is about $\lambda_{\text{cr}}/2$ (Salo et al. 2004).

After reaching about unity Q increases with time and finally the time-averaged Q becomes almost constant $\simeq 2$ after 5 orbital periods. In this state, the radial spacing is about λ_{cr} (Daisaka & Ida 1999; Salo et al. 2004). Since heating by gravitational scattering and cooling by inelastic collisions are balanced, the time-averaged velocity dispersion is constant. Figure 1c shows the equilibrium state of the self-gravity wakes. The high density region is trailing and inclined by 10–30° with respect to the tangential direction.

We performed simulations with various ring parameters and confirmed that the equilibrium state is reached after 5 orbital periods in all models if the axisymmetric overstability does not exist. Thus, hereafter we analyze the simulation result only for $\tilde{t} = 5$ –20.

3.2. Two-Dimensional Autocorrelation Function

Since the self-gravity wakes are a transient and recurrent feature we need to investigate them in a statistical way. Two-dimensional autocorrelation function has been widely used to analyze self-gravity wakes (Toomre & Kalnajs 1991; Salo 1995; Daisaka & Ida 1999; Salo et al. 2004). We use the two-dimensional autocorrelation function of the surface density given by

$$\xi(x, y, t) = \frac{1}{\Sigma_0^2 L^2} \int_{-L/2}^{L/2} \int_{-L/2}^{L/2} \Sigma(x + x', y + y', t) \Sigma(x', y', t) dx' dy' - 1. \quad (11)$$

Table 1—Continued

Model	τ	$\tilde{r}_H$	ϵ	N	Σ_0 (g cm^{-2})	a (10^5 km)	L (km)	θ_i (deg.)	θ_o (deg.)	x_b (λ_{cr})	$ y_b $ (λ_{cr})
41	0.60	0.98	0.50	138497	36–72	0.95–1.20	0.85	39.0 ± 7.6	14.0 ± 7.4	0.31	0.38
42	0.60	1.02	0.50	176070	36–72	0.99–1.24	0.96	41.9 ± 7.2	14.7 ± 6.8	0.34	0.37
43	0.60	1.06	0.50	221779	36–72	1.03–1.29	1.08	46.4 ± 8.9	6.7 ± 6.1	0.56	0.53
44	0.90	0.68	0.50	52169	54–108	0.66–0.83	0.43	15.0 ± 1.7	13.4 ± 1.8	0.21	0.78
45	0.90	0.73	0.50	79854	54–108	0.71–0.89	0.53	21.6 ± 2.1	14.8 ± 3.1	0.09	0.22
46	0.90	0.77	0.50	109977	54–108	0.75–0.94	0.62	22.6 ± 3.2	14.2 ± 5.3	0.19	0.45
47	0.90	0.81	0.50	149028	54–108	0.78–0.99	0.72	24.5 ± 3.9	12.8 ± 5.8	0.35	0.76
48	0.90	0.85	0.50	199009	54–108	0.82–1.04	0.83	26.2 ± 6.1	17.7 ± 7.1	0.28	0.57
49	0.90	0.90	0.50	280423	54–108	0.87–1.10	0.99	31.2 ± 5.2	14.7 ± 5.8	0.30	0.49
50	0.90	0.94	0.50	364021	54–108	0.91–1.15	1.13	33.0 ± 9.9	9.8 ± 7.6	0.49	0.76
51	0.90	0.98	0.50	467429	54–108	0.95–1.20	1.28	39.0 ± 8.9	14.2 ± 5.7	0.28	0.35
52	0.90	1.02	0.50	594238	54–108	0.99–1.24	1.44	38.8 ± 11.6	11.7 ± 9.4	0.56	0.70
53	0.90	1.06	0.50	748505	54–108	1.03–1.29	1.62	47.4 ± 10.4	14.3 ± 5.6	0.29	0.27




Fig. 1.— Snapshots of the surface density normalized by the mean surface density Σ_0 at $\tilde{t} = 0.0$ (left), $\tilde{t} = 1.0$ (middle) and $\tilde{t} = 5.0$ (right) for model 1.

Table 2. Ring Parameters and Pitch Angles for Size Distribution Models

Model	a (10^5 km)	ρ_p (g cm^{-3})	$R_{\min}-R_{\max}$ (cm)	N	L (km)	θ_i (deg.)	θ_o (deg.)	x_b (λ_{cr})	$ y_b $ (λ_{cr})
54	1.30	0.45	150–150	46148	0.76	30.0 ± 5.0	14.8 ± 8.3	0.21	0.36
55	1.30	0.45	106–212	47421	0.76	31.6 ± 8.5	15.6 ± 6.9	0.18	0.30
56	1.30	0.45	87–260	49392	0.76	34.3 ± 8.0	14.8 ± 6.9	0.20	0.30
57	1.30	0.45	75–300	52260	0.76	34.1 ± 8.0	13.7 ± 6.4	0.26	0.39
58	1.30	0.45	67–335	54745	0.76	35.5 ± 5.6	13.4 ± 5.3	0.27	0.38
59	1.30	0.45	61–367	57128	0.76	37.5 ± 5.6	14.9 ± 7.0	0.23	0.30
60	1.30	0.45	57–397	58069	0.76	41.1 ± 8.5	15.8 ± 5.9	0.21	0.24
61	1.30	0.45	53–424	60591	0.76	40.9 ± 7.2	14.7 ± 5.8	0.24	0.28
62	1.30	0.45	50–450	62207	0.76	41.1 ± 6.9	15.1 ± 6.1	0.26	0.29
63	1.30	0.45	47–474	64929	0.76	41.6 ± 5.0	15.1 ± 4.7	0.25	0.28
64	0.85	0.45	47–474	4998	0.21	22.0 ± 8.9	18.2 ± 5.5	0.67	1.66
65	0.90	0.45	47–474	7043	0.25	28.0 ± 5.1	11.0 ± 4.2	1.02	1.92
66	0.95	0.45	47–474	9742	0.30	22.6 ± 6.4	18.8 ± 3.9	0.36	0.87
67	1.00	0.45	47–474	13254	0.35	34.2 ± 4.7	15.2 ± 4.0	0.26	0.39
68	1.05	0.45	47–474	17761	0.40	30.4 ± 5.7	12.8 ± 5.2	0.41	0.70
69	1.10	0.45	47–474	23480	0.46	31.4 ± 4.3	10.7 ± 5.7	0.50	0.82
70	1.15	0.45	47–474	30657	0.53	32.8 ± 6.5	13.7 ± 7.1	0.30	0.47
71	1.20	0.45	47–474	39576	0.60	34.8 ± 5.3	12.5 ± 6.6	0.32	0.46
72	1.25	0.45	47–474	50560	0.68	35.9 ± 7.1	14.4 ± 5.4	0.29	0.40
73	1.30	0.45	47–474	63974	0.76	43.0 ± 5.7	14.6 ± 5.2	0.25	0.26
74	1.35	0.45	47–474	80232	0.85	41.1 ± 8.2	8.8 ± 6.4	0.48	0.55
75	0.85	0.90	47–474	2499	0.21	24.8 ± 13.9	19.4 ± 8.5	0.89	1.92
76	0.90	0.90	47–474	3521	0.25	36.6 ± 11.0	16.0 ± 8.5	0.53	0.72
77	0.95	0.90	47–474	4871	0.30	37.9 ± 21.0	14.8 ± 10.4	0.34	0.44
78	1.00	0.90	47–474	6627	0.35	44.3 ± 16.2	13.9 ± 8.0	0.48	0.49
79	1.05	0.90	47–474	8880	0.40	44.9 ± 13.4	8.6 ± 7.9	0.93	0.93
80	1.10	0.90	47–474	11740	0.46	44.3 ± 10.1	15.4 ± 9.2	0.42	0.43
81	1.15	0.90	47–474	15328	0.53	45.6 ± 26.1	12.5 ± 9.1	0.49	0.48
82	1.20	0.90	47–474	19788	0.60	54.9 ± 7.0	12.6 ± 8.2	0.48	0.33
83	1.25	0.90	47–474	25280	0.68	58.0 ± 8.0	12.8 ± 6.7	0.51	0.32
84	1.30	0.90	47–474	31987	0.76	90.4 ± 8.8	16.3 ± 11.2	0.22	0.00
85	1.35	0.90	47–474	40116	0.85	90.2 ± 2.9	16.2 ± 8.7	0.28	0.00

Note. — The surface density is $\Sigma_0 = 50 \text{ g cm}^{-2}$ and the restitution coefficient is $\epsilon = 0.5$.




Fig. 2.— Time evolution of Q for model 1.

We calculate the autocorrelation with the uniform grid of 256×256 . We first calculate the surface density at each grid from particle distribution and then evaluate the integral. Examples are shown in Figure 3. Though there are particle aggregates with various shapes in the surface density snapshot, the autocorrelation analysis enables us to extract a smooth wake structure. As shown in Figure 3a, b, and c, although the wake structure is almost steady, it slightly fluctuates with time. Therefore we introduce the time-averaged autocorrelation:

$$\bar{\xi}(x, y) = \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} \xi(x, y, t') dt', \quad (12)$$

where we set $\tilde{t}_1 = 5$ and $\tilde{t}_2 = 20$. We approximate the time integral by the sum of 150 snapshots in $\tilde{t} = 5.0\text{--}20.0$ with $\Delta\tilde{t} = 0.1$. Figure 3d shows the time-averaged autocorrelation over $\tilde{t} = 5.0\text{--}20.0$. The small variation is completely smoothed out. The inclined structure at the center corresponds to the structure of the high density region in the self-gravity wakes. This inclination means that the structure is trailing and the pitch angle is about 20° . The regions with negative autocorrelation coefficient correspond to voids between self-gravity wakes.

It should be noted that the grid of 256×256 is reasonable. One might worry that the number of grids larger than that of particles leads to large density error. This is always true for low density parts where particles are sparsely distributed even if the number of particles are larger than that of grids. However, it is not problematic here since we focus on high density parts automatically considering the autocorrelation function. In addition, time-averaging of the autocorrelation function reduces the density error. To demonstrate a suitable grid size Figure 4 demonstrates the validity of the grid size. The function shape is correctly caught by the grid of 256×256 . The typical discretization error of the pitch angle is smaller than $\simeq 4\%$.

3.3. Calculation of Pitch Angle

We calculate the pitch angle from the time-averaged autocorrelation. First, we extract the ridge of the dense part of autocorrelation by considering the integral curve of the gradient field of the autocorrelation function. Figure 5 shows examples of the ridge of autocorrelation, which confirms the previous results that the slope of the ridge line varies with the distance from the center (Salo et al. 2004). The slope in the outer part of the autocorrelation function is steeper than that in the inner part. Clearly we cannot fit the ridge by one line. Therefore we fit the ridge with the following function:

$$y = -\cot \theta_o x + (\cot \theta_o - \cot \theta_i) x_b \operatorname{erf} \left(\frac{\sqrt{\pi} x}{2x_b} \right), \quad (13)$$

where θ_i , θ_o , and x_b are parameters. The slope of the tangent to the fitting curve around the center $|x| \ll x_b$ is $\cot \theta_i$ and that for $|x| \gg x_b$ is $\cot \theta_o$. Hereafter, we define the inner and outer pitch angles as θ_i and θ_o , where θ_i and θ_o correspond to the pitch angles in the inner and outer parts of




Fig. 3.— Snapshots of the two-dimensional autocorrelation function at $\tilde{t} = 5.0$ (a), 10.0 (b), and 15.0 (c) for model 1. The panel (d) shows the time-averaged autocorrelation over $\tilde{t} = 5.0$ –20.0.

the autocorrelation function, respectively. The value x_b means the typical length where the pitch angle changes from θ_i to θ_o . An example of the fitting function is shown in Figure 6.

We introduce a line integral of autocorrelation function along the ridge defined by Equation (13) as

$$S(\theta_i, \theta_o, x_b) = \int \bar{\xi}(x, y) dl. \quad (14)$$

We calculate the integral by Simpson’s rule and find θ_i , θ_o , and x_b that maximize S by the downhill simplex method (Nelder & Mead 1965; Press 2007).

We obtain the representative pitch angles from the time averaged autocorrelation function. However, the actual pitch angles fluctuate with time around the representative values. Thus, in order to estimate the fluctuation degree, we calculate the median absolute deviation from the time series of the inner and outer pitch angles calculated from the autocorrelation functions for $\tilde{t} = 5.0\text{--}20.0$. Examples of the optimized parameters and the median absolute deviation are $\theta_i = (25.8 \pm 4.1)^\circ$, $\theta_o = (13.3 \pm 4.2)^\circ$, and $x_b/\lambda_{cr} = 0.26$ for model 1 and $\theta_i = (41.9 \pm 7.2)^\circ$, $\theta_o = (14.7 \pm 6.8)^\circ$, and $x_b/\lambda_{cr} = 0.34$ for model 42. As shown in Figure 5, the ridge curves agree well with Equation (13). Model 42 shows the deviation of the extracted ridge from the fitting curve in the low autocorrelation region. This is because that the ridge extracted by the gradient field is sensitively affected by the noise. In the low autocorrelation region, the fitting curve is more robustly determined than the extracted ridge.

The inner and outer pitch angles θ_i and θ_o , and the breakpoint position (x_b, y_b) are summarized in Tables 1 and 2 where y_b is calculated from $y_b = -x_b/\tan\theta_i$. The breakpoint x_b is generally less than λ_{cr} and about $(0.1\text{--}0.5)\lambda_{cr}$.

4. Parameter Dependence

We first adopt the identical particle model and examine the dependence of the pitch angle on the restitution coefficient ϵ , the optical depth τ , and the Hill radius of a particle pair relative to the sum of their physical radii $\tilde{r}_H$. Next, we investigate the effect of the size distribution of ring particles.

4.1. Restitution Coefficient

First we adopt the constant restitution coefficient model and investigate its effect on the pitch angle. The restitution coefficients adopted here are $\epsilon = 0.1, 0.2, 0.3, 0.4, 0.5$ and 0.6 (models 2–7). The other parameters are $\tau = 0.7$ and $\tilde{r}_H = 0.81$. The results are summarized in Figure 7. In all models, the inner and outer pitch angles barely depend on ϵ , and their mean values are about 26.1° and 14.1° , respectively. As shown by Salo et al. (2004), the inner pitch angle is larger than the outer one. Since the influence of the self-gravity relative to the shear is strong around the center of




Fig. 4.— Time-averaged autocorrelation functions for model 30 with grid of (a) 64×64 , (b) 128×128 , and (c) 256×256 .




Fig. 5.— Autocorrelation function and the extracted ridge (dotted) for models 1 (left) and 42 (right). The dashed curve shows the fitting function given by Equation (13) .




Fig. 6.— Fitting function for the ridge of autocorrelation with $\theta_i = 50^\circ$, $\theta_o = 20^\circ$, and $x_b = 1$ (solid line). The dotted line denotes the asymptotic lines the pitch angles of which are θ_i and θ_b . The breakpoint is x_b .

self-gravity wakes, the shape in the inner part of the autocorrelation function is nearly spherical. Conversely, the shape in the outer part is elongated due to the shear. Thus, we have $\theta_i > \theta_o$.

It is expected that the pitch angle does not change for the velocity-dependent restitution coefficient model, since the pitch angles barely depend on ϵ in the constant restitution coefficient models. This is confirmed by the simulation with the velocity-dependent restitution coefficient (model 8) where we assume $a = 1.25 \times 10^5$ km, $\rho_p = 0.45$ g cm $^{-3}$, and $R_p = 120$ cm (Fig.7). The inner and outer pitch angles are 24.5° and 13.9° , respectively. They are approximately the same as those for the constant restitution coefficient models. It is clearly shown that the pitch angle is independent of the restitution coefficient.

4.2. Optical Depth

We investigate the dependence of the pitch angle on the dynamical optical depth τ with fixed $\tilde{r}_H$ and ϵ . The restitution coefficient is $\epsilon = 0.5$ and the Hill radius of a particle pair relative to the sum of their physical radii is $\tilde{r}_H = 0.6$ and 0.81 . The optical depths are $\tau = 0.8$ – 1.8 for $\tilde{r}_H = 0.6$ (models 9–14) and $\tau = 0.3$ – 1.3 for $\tilde{r}_H = 0.81$ (models 15–24). The results are shown in Figure 8. For $\tilde{r}_H = 0.6$ the inner pitch angle roughly decreases with τ , and becomes lower than 10° . For $\tau > 1.0$ the outer pitch angle is larger than the inner one. However, it is arguable whether these values can be considered as the representative pitch angle of self-gravity wakes. Figure 9 illustrates the surface density and its autocorrelation for $\tau = 1.6$ and $\tilde{r}_H = 0.6$ (model 13) at $\tilde{t} = 19$. We find that the axisymmetric structure and the non-axisymmetric self-gravity wakes coexist. This axisymmetric structure is caused by the overstable oscillation (Schmit & Tscharnuter 1995; Schmidt et al. 2001; Salo et al. 2001; Schmidt & Salo 2003; Latter & Ogilvie 2006). The axisymmetric structure appears for $\tau \gtrsim 1$ and $\tilde{r}_H \sim 0.6$. The overstable oscillation causes the small pitch angles (French et al. 2007).

On the other hand, for $\tilde{r}_H = 0.81$ the outer pitch angle does not depend on τ , and for $\tau > 0.5$ the inner pitch angle does not have clear dependence on τ . For $\tau > 0.5$ the mean inner and outer pitch angles are 24.7° and 13.7° , respectively. For $0.3 \leq \tau \leq 0.5$ the inner pitch angle is systematically larger than 24.7° . The inner pitch angles are 30.2° , 30.3° , and 29.5° for $\tau = 0.3$ (model 15), 0.4 (model 16), and 0.5 (model 17), respectively. Figure 10 represents the surface density snapshots at $\tilde{t} = 20$ for $\tau = 0.30$ (model 15), 0.40 (model 16), 0.50 (model 17), and 0.60 (model 18). From Equation (10), the condition for self-gravity wake formation is $\tau \gtrsim 0.27$. Thus the ring of model 15 is marginally unstable and the self-gravity wakes are small and faint. As τ increases (models 16, 17, and 18), the clearer long self-gravity wakes and voids develop. A possible explanation for the large inner pitch angle at small τ is as follows. In marginally unstable rings, the self-gravity is weak and the resulting self-gravity wakes are faint. Since the wake clumps are loosely bound, they are easily destroyed by the interactions among clumps before they are elongated. In short we find that as long as the overstable oscillation does not occur and the clear self-gravity wakes form with large τ , the inner pitch angle is almost independent of τ .




Fig. 7.— Inner and outer pitch angles θ_i and θ_o are plotted against the restitution coefficient ϵ with $\tau = 0.7$ and $\tilde{r}_H = 0.81$ (models 2–7). The open and filled symbols denote θ_i and θ_o , respectively. The solid and dashed lines denote θ_i and θ_o for the velocity-dependent restitution coefficient model, respectively (model 8).




Fig. 8.— Inner and outer pitch angles θ_i (open) and θ_o (filled) are plotted against the optical depth τ . The restitution coefficient is $\epsilon = 0.5$ and Hill radius of a particle pair relative to the sum of their physical radii is $\tilde{\tau}_H = 0.6$ (left, models 9–14) and 0.81 (right, models 15–24).




Fig. 9.— Snapshots of the surface density (left) and its autocorrelation (right) for the model with $\tau = 1.6$, $\tilde{\tau}_H = 0.6$, and $\epsilon = 0.5$ (model 15) at $\tilde{t} = 19.0$.




Fig. 10.— Snapshots of the surface density normalized by the mean surface density at $\tilde{t} = 20.0$ for models 15 (a), 16 (b), 17 (c), and 18 (d).

4.3. Hill Radius Relative to Sum of Particles Radii

The dependence of the pitch angle on $\tilde{r}_H$ is shown for $\tau = 0.3$ (models 25–34), 0.6 (models 35–43), and 0.9 (models 44–53) in Figure 11. The outer pitch angle is independent of both τ and $\tilde{r}_H$. The mean outer pitch angles for $\tau = 0.3, 0.6$, and 0.9 are 12.0° , 12.8° , and 13.8° , respectively. This reason will be discussed in Section 5.1.

On the other hand, the inner pitch angle increases with $\tilde{r}_H$ from 30° to 50° for $\tilde{r}_H > 0.8$. The similar tendency was shown in the previous numerical studies (Salo 1995; Takeda & Ida 2001). However, for $\tilde{r}_H < 0.8$ and $\tau = 0.6$ (models 35, 36, and 37), the inner pitch angle is almost constant, and for $\tilde{r}_H = 0.77$ and $\tau = 0.3$ (model 27) the inner pitch angle is unexpectedly large. The rings of these models are marginally unstable. Thus, as discussed in Section 4.2, the pitch angle becomes large.

For $0.8 < \tilde{r}_H$, using the least-square fit method, we obtain the fitting formula of the inner pitch angle:

$$\theta_i \simeq \begin{cases} (117.9 \pm 12.0)\tilde{r}_H + (-65.1 \pm 11.3) \text{ deg.} & (\tau = 0.3), \\ (94.8 \pm 4.9)\tilde{r}_H + (-52.0 \pm 4.6) \text{ deg.} & (\tau = 0.6), \\ (113.3 \pm 21.6)\tilde{r}_H + (-69.5 \pm 20.4) \text{ deg.} & (\tau = 0.9), \end{cases} \quad (15)$$

which shows that the dependence of θ_i on $\tilde{r}_H$ is independent of τ . Thus, fitting the results of all τ models we obtain a general formula for $\tilde{r}_H > 0.8$ and $\tau = 0.3$ –0.9:

$$\theta_i \simeq (108.7 \pm 13.4)\tilde{r}_H + (-62.2 \pm 12.8) \text{ deg.} \quad (16)$$

4.4. Size Distribution

We investigate the effect of the size distribution of ring particles on the pitch angle. We adopt the power-law size distribution described by Equation (6) with the lower and upper size limits




Fig. 11.— Inner and outer pitch angles θ_i (open) and θ_o (filled) plotted against $\tilde{r}_H$ with the optical depth $\tau = 0.3$ (left, models 25–34), $\tau = 0.6$ (middle, models 35–43), and $\tau = 0.9$ (right, models 44–53). The solid lines denote the fitting formula given by Equations (16), and the dashed lines denote the estimation given by Equation (22) with $C_w = 0.45$ and $C_d = 1.0$.

(models 54–63). Figure 12 illustrates the dependence of the pitch angle on the ratio $R_{\max}/R_{\min}$. The surface density, the Saturnicentric distance, and the particle density are $\Sigma_0 = 50 \text{ g cm}^{-2}$, $a = 1.3 \times 10^5 \text{ km}$, and $\rho_p = 0.45 \text{ g cm}^{-3}$, respectively. The inner pitch angle increases with $R_{\max}/R_{\min}$. The inner pitch angle with the size distribution is 30.0° – 41.6° and is 39% larger than that for identical particles. This is consistent with the previous results by Salo et al. (2004). As $R_{\max}/R_{\min}$ increases, the filling factor and density of self-gravity wakes increase. The higher density of self-gravity wakes leads to the larger pitch angle because the shear in the higher density wakes is weaker relative to the self-gravity. The details of this dependence will be discussed in Section 5.2. On the other hand, the outer pitch angle barely depends on $R_{\max}/R_{\min}$. The mean outer pitch angle is 14.8° .

In Section 4.3, we found that the pitch angle for identical particles increases with $\tilde{r}_H$. We expect the same trend for the pitch angle in the size distribution models. Figure 13 shows the dependence of the pitch angle on the Saturnicentric distance. The surface density is $\Sigma_0 = 50 \text{ g cm}^{-2}$ and the particle density is $\rho_p = 0.45 \text{ g cm}^{-3}$ (models 63–73) and 0.9 g cm^{-3} (models 74–84). The lower and upper size limits are fixed as $R_{\min} = 41.7 \text{ cm}$ and $R_{\max} = 417 \text{ cm}$. The inner pitch angles for $\rho_p = 0.45 \text{ g cm}^{-3}$ are smaller than those for $\rho_p = 0.9 \text{ g cm}^{-3}$. This is because smaller ρ_p corresponds to smaller $\tilde{r}_H$. As expected, the inner pitch angle generally increases with the Saturnicentric distance. For $\rho_p = 0.45 \text{ g cm}^{-3}$ the pitch angle increases with the Saturnicentric distance a from 22.0° to 43.0° , and the outer pitch angle has no trend and ranges within 8.8° – 18.8° . Similarly, for $\rho_p = 0.9 \text{ g cm}^{-3}$, the pitch angle increases with a from 24.8° to 58.0° for $a \leq 1.25 \times 10^5 \text{ km}$, and the outer pitch angle is 8.6° – 19.4° .

For $\rho_p = 0.9 \text{ g cm}^{-3}$ and $a \geq 1.3 \times 10^5 \text{ km}$, the inner pitch angle is about 90° . Figure 14 presents the snapshot of the surface density and the autocorrelation for $a = 1.35 \times 10^5 \text{ km}$ and $\rho_p = 0.9 \text{ g cm}^{-3}$ (model 84). In the surface density snapshot, we cannot observe the distinct elongated wakes but nearly spherical aggregates. This is because the aggregate formation starts from $a \simeq 1.3 \times 10^5 \text{ km}$ for $\rho_p = 0.9 \text{ g cm}^{-3}$ (Ohtsuki 1993; Salo 1995; Karjalainen & Salo 2004). In the autocorrelation function, the shape of the dense part is an ellipsoid, which corresponds to the Hill sphere. The long axis of the ellipsoid is almost parallel to the x -axis. Thus, in this case, the inner pitch angle becomes around 90° . The typical mass of the self-gravity wake is $\sim \Sigma_0 \lambda_{\text{cr}}^2$. Thus, supposing that a particle with mass $\Sigma_0 \lambda_{\text{cr}}^2$ is located at the center, we obtain the corresponding Hill radius $(\Sigma_0 \lambda_{\text{cr}}^2 / 3M_S)^{1/3} a = \lambda_{\text{cr}} / (12\pi^2)^{1/3} \simeq 0.2 \lambda_{\text{cr}}$. This length scale is consistent with the size of the ellipsoid in autocorrelation function.

We compare the simulation results with the observations. Hedman et al. (2007) found that the pitch angle increases from 17° to 26° with the Saturnicentric distance in the A ring. Colwell et al. (2007) found the similar results that the pitch angle in the A ring is 10 – 50° and increases with the Saturnicentric distance. Our simulations with $\rho_p = 0.45 \text{ g cm}^{-3}$ show that the inner pitch angle increases with the Saturnicentric distance around $a \simeq 1.3 \times 10^5 \text{ km}$ and the inner and outer pitch angles are 43.0° and 14.6° . We expect that in general the observed pitch angle ranges within the inner and outer pitch angles. Therefore, our simulations with $\rho_p = 0.45 \text{ g cm}^{-3}$ are consistent




Fig. 12.— Inner and outer pitch angles θ_i (open) and θ_o (filled) plotted against $R_{\max}/R_{\min}$ for models 54–63. The surface density, the Saturnicentric distance and the particle density are $\Sigma_0 = 50 \text{ g cm}^{-2}$, $a = 1.3 \times 10^5 \text{ km}$ and $\rho_p = 0.45 \text{ g cm}^{-3}$, respectively. The solid and dashed lines denote the estimation given by Equation (22) with $C_w = 0.45$ (solid) and $C_w = 1.0$ (dashed), respectively.




Fig. 13.— Inner and outer pitch angles θ_i (open) and θ_o (filled) plotted against the Saturnicentric distance. The particle densities are $\rho_p = 0.45 \text{ g cm}^{-3}$ (left, models 63–73) and $\rho_p = 0.9 \text{ g cm}^{-3}$ (right, models 74–84). The solid lines denote the estimation given by Equation (22) with $C_w = 1.0$ and $C_d = 1.0$.




Fig. 14.— Surface density snapshot at $\tilde{t} = 20.0$ (left) and its time-averaged autocorrelation (right) with the extracted ridge (dotted) and the fitting function (dashed) for model 84.

with the observations.

5. Discussion

5.1. Comparison with Galactic Spirals

We compare the pitch angles of spiral arms in disk galaxies and self-gravity wakes in planetary rings. Disk galaxies are considered as collisionless systems. Recently, the pitch angle of spiral arms was investigated by the numerical and analytic studies (Grand et al. 2013; Michikoshi & Kokubo 2014). In the galactic disk, the pitch angle depends only on the shear rate defined by

$$\Gamma = 2 - \frac{\kappa^2}{2\Omega^2}, \quad (17)$$

where κ is the epicycle frequency. From the numerical simulations and the linear analysis Michikoshi & Kokubo (2014) obtained the fitting formula of the pitch angle

$$\tan \theta = \frac{2}{7} \frac{\sqrt{4 - 2\Gamma}}{\Gamma}. \quad (18)$$

For the shear rate of the Keplerian rotation $\Gamma = 3/2$, we obtain the pitch angle $\theta \simeq 11^\circ$, which is consistent with the outer pitch angle of self-gravity wakes.

In the central dense parts where self-gravity wakes are closely packed, inelastic collisions play an important role in dynamics. Thus, the inner pitch angle of self-gravity wakes deviates from the estimated value for collisionless systems. In the outer low density regions, however, the effect of the shear is relatively strong compared to that of collisions, and the shear rate mainly determines the outer pitch angle just like collisionless systems. Thus, the outer pitch angle is the same as that of collisionless systems and is independent of any ring parameters.

5.2. Inner Pitch Angle

We simplify the physical processes concerning the inner pitch angle and derive a rough estimate as follows. In high density regions, the self-gravity is strong. Without the shear and only with the self-gravity, an ellipsoidal aggregate whose shape is the Hill sphere would form and its long axis is parallel to x -axis, that is, the pitch angle is 90° . Therefore, the strong self-gravity tends to increase the inner pitch angle to 90° . Conversely, the shear rotates the aggregate and tends to decrease the inner pitch angle to 0° . The inner pitch angle may be determined by the balance between these two processes.

We consider the evolution of an aggregate formed due to the self-gravity. When the aggregate is deformed by the shear, the self-gravity tends to restore it back to its original shape. The restoration timescale may be approximated by $t_{\text{grav}} \sim 1/\sqrt{G\rho_w}$ where ρ_w is the bulk density of

self-gravity wakes, which is the timescale of the Jeans instability or gravitational instability (e.g., Chandrasekhar & Fermi 1953). The bulk density ρ_w is less than the particle density ρ_p . Introducing a dimensionless factor C_w , we assume $\rho_w = C_w \rho_p$. For identical particles, C_w is less than 0.74 that is for the closest packing. The timescale of self-gravity is given by

$$t_{\text{grav}} \simeq \frac{C_d}{\sqrt{GC_w \rho_p}} = \frac{C_d}{3\Omega} \sqrt{\frac{\pi}{C_w}} \tilde{r}_H^{-3/2}, \quad (19)$$

where C_d is a non-dimensional factor on the order of unity. The shear rotates and extends the wake and the pitch angle decreases with t as

$$\tan \theta = \frac{2}{3\Omega t}, \quad (20)$$

where t is the time elapsed since $\theta = 90^\circ$ (e.g., Toomre 1981; Binney & Tremaine 2008). The shear timescale is

$$t_{\text{shear}} \simeq \left| \frac{1}{\tan \theta} \frac{d \tan \theta}{dt} \right|^{-1} = \frac{2}{3\Omega \tan \theta}. \quad (21)$$

As the pitch angle decreases with time, the wake rotation slows down.

When the wake pitch angle is large ($\theta \sim 90^\circ$), the wake rotation is fast ($t_{\text{shear}} < t_{\text{grav}}$). The wake rotates quickly before the self-gravity restores the inclined wake back to its original shape. As the pitch angle decreases, the wake rotation slows down and the self-gravity becomes important. Thus, from $t_{\text{shear}} = t_{\text{grav}}$ we obtain a crude estimate of the inner pitch angle:

$$\tan \theta_i \simeq \frac{2}{C_d} \sqrt{\frac{C_w}{\pi}} \tilde{r}_H^{3/2} = 0.84 \frac{\sqrt{C_w}}{C_d} \left(\frac{\rho_p}{0.9 \text{ g cm}^{-3}} \right)^{1/2} \left(\frac{a}{10^5 \text{ km}} \right)^{3/2}, \quad (22)$$

which means that the pitch angle increases with C_w . The bulk density of self-gravity wakes with size distribution can be larger than that for identical particles, and thus the pitch angle with size distribution is larger than that for identical particles, which is consistent with the simulation results. Figure 12 presents the pitch angle as a function of $R_{\text{max}}/R_{\text{min}}$. As $R_{\text{max}}/R_{\text{min}}$ increases, the filling factor C_w increases. Assuming $C_d = 1.0$, we find that $C_w = 0.45$ and 1 are suitable for the identical particle models and the size distribution models of $R_{\text{max}}/R_{\text{min}} = 10$, respectively. As shown in Figures 11 and 13, Equation (22) can roughly explain the dependence of the inner pitch angle on $\tilde{r}_H$ and a with $C_w = 0.45$ for the identical particle models and $C_w = 1.0$ for the size distribution models with $R_{\text{max}}/R_{\text{min}} = 10$. The simulation results and Equation (22) show that the inner pitch angle increases with $\tilde{r}_H$. Since $\tilde{r}_H$ is proportional to the Saturnicentric distance a , as $\tilde{r}_H$ increases, the orbital period increases, that is, the shear timescale increases. On the other hand, the timescale of the gravitational instability depends only on ρ_w and is independent of a . Therefore, as $\tilde{r}_H$ increases, the self-gravity dominates over the shear and the inner pitch angle increases.

Note that the actual filling factor has not been investigated in detail yet. These results do not suggest that the filling factor is $C_w = 0.45$ or 1.0. The factor C_d is also a free parameter. If

we adopt smaller C_d , the corresponding filling factor becomes smaller. The realistic filling factor should be smaller than unity.

We used $R_{\max}/R_{\min} \leq 10$ in the size distribution models. However, this size ratio is far smaller than the observationally inferred values that are 70 and 2000 in the inner and outer A rings, respectively (French & Nicholson 2000). As shown in Figure 12, the inner pitch angle is almost constant for $R_{\max}/R_{\min} > 5$. Accordingly, the simulation results with $R_{\max}/R_{\min} = 10$ in Section 4.4 may be applicable to the realistic ring.

6. Summary

We have investigated the pitch angle of the self-gravity wakes in the dense planetary ring by performing local N -body simulations where both inelastic collisions and self-gravitational interactions between particles are included. In order to extract the feature of the self-gravity wakes, we calculated the two-dimensional autocorrelation function of the surface density. We measured the angle of the inclined structure of the autocorrelation function as the wake pitch angle. We obtained two pitch angles in the inner and outer parts of the autocorrelation function separately. We confirmed that the inner pitch angle is larger than the outer one in all models.

We examined the dependence of the pitch angle on various ring parameters. We found that the outer pitch angle is independent of the ring parameters, which is 10° – 15° . This value is consistent with the estimation of pitch angle of the spiral arms in collisionless systems (Julian & Toomre 1966; Michikoshi & Kokubo 2014). The inner pitch angle does not depend on the restitution coefficient ϵ of particles and the optical depth τ provided that the overstable oscillation does not occur. We found that the inner pitch angle is determined by the Hill radius of a particle pair relative to the sum of their physical radii $\tilde{r}_H$. The inner pitch angle increases with $\tilde{r}_H$ from 20° to 50° . The tendency that the pitch angle increases with $\tilde{r}_H$ is consistent with the result of the observational studies that the pitch angle increases with the Saturn-centric distance (Hedman et al. 2007; Colwell et al. 2007). The pitch angle with the size distribution model is larger than that for the identical particles because the filling factor for the size distribution model is larger and the resulting self-gravity is stronger. The strong self-gravity increases the inner pitch angle.

Considering the effects of the self-gravity and the shear, the tendency of the increase of the inner pitch angle with the Saturn-centric distance can be explained. However, this model is far from a complete theory. We need a more sophisticated theoretical framework to understand the overall activity of the self-gravity wakes and more detailed analyses of the self-gravity wakes by N -body simulation. It is especially important to understand the wake amplification and destruction processes considering the inelastic collisions and the nonlinear effects. We will focus on these processes in the next paper.

Numerical computations were carried out on the GRAPE system at Center for Computational

Astrophysics, National Astronomical Observatory of Japan.

REFERENCES

- Binney, J. & Tremaine, S. 2008, *Galactic dynamics*, (Princeton university press)
- Bridges, F. G., Hatzes, A., & Lin, D. N. C. 1984, *Nature*, 309, 333
- Camichel, H. 1958, *Annales d’Astrophysique*, 21, 231
- Chandrasekhar, S. & Fermi, E. 1953, *ApJ*, 118, 116
- Colombo, G., Goldreich, P., & Harris, A. W. 1976, *Nature*, 264, 344
- Colwell, J. E., Esposito, L. W., Sremčević, M., Stewart, G. R., & McClintock, W. E. 2007, *Icarus*, 190, 127
- Daisaka, H. & Ida, S. 1999, *Earth, Planets, and Space*, 51, 1195
- Daisaka, H., Tanaka, H., & Ida, S. 2001, *Icarus*, 154, 296
- Dones, L., Cuzzi, J. N., & Showalter, M. R. 1993, *Icarus*, 105, 184
- Ferrin, I. 1975, *Ap&SS*, 33, 453
- Franklin, F. A., Cook, A. F., Barrey, R. T. F., Roff, C. A., Hunt, G. E., & de Rueda, H. B. 1987, *Icarus*, 69, 280
- French, R. G. & Nicholson, P. D. 2000, *Icarus*, 145, 502
- French, R. G., Salo, H., McGhee, C. A., & Dones, L. 2007, *Icarus*, 189, 493
- Goldreich, P. & Tremaine, S. D. 1978, *Icarus*, 34, 227
- Grand, R. J. J., Kawata, D., & Cropper, M. 2013, *A&A*, 553, A77
- Hedman, M. M., Nicholson, P. D., Salo, H., Wallis, B. D., Buratti, B. J., Baines, K. H., Brown, R. H., & Clark, R. N. 2007, *AJ*, 133, 2624
- Julian, W. H. & Toomre, A. 1966, *ApJ*, 146, 810
- Karjalainen, R. & Salo, H. 2004, *Icarus*, 172, 328
- Latter, H. N. & Ogilvie, G. I. 2006, *MNRAS*, 372, 1829
- Makino, J., Hiraki, K., & Inaba, M. 2007, *Proceedings of the 2007 ACM/IEEE conference*, 1
- Michikoshi, S. & Kokubo, E. 2011, *ApJ*, 732, L23+

- . 2014, *ApJ*, 787, 174
- Nakazawa, K. & Ida, S. 1988, *Progress of Theoretical Physics Supplement*, 96, 167
- Nelder, J. A. & Mead, R. 1965, *The computer journal*, 7, 308
- Ohtsuki, K. 1993, *Icarus*, 106, 228
- Ohtsuki, K. & Emori, H. 2000, *AJ*, 119, 403
- Petit, J.-M. & Henon, M. 1986, *Icarus*, 66, 536
- Porco, C. C., Weiss, J. W., Richardson, D. C., Dones, L., Quinn, T., & Throop, H. 2008, *AJ*, 136, 2172
- Press, W. H. 2007, *Numerical recipes 3rd edition: The art of scientific computing*, (Cambridge university press)
- Reitsema, H. J., Beebe, R. F., & Smith, B. A. 1976, *AJ*, 81, 209
- Richardson, D. C. 1994, *MNRAS*, 269, 493
- Robbins, S. J., Stewart, G. R., Lewis, M. C., Colwell, J. E., & Sremčević, M. 2010, *Icarus*, 206, 431
- Salo, H. 1991, *Icarus*, 90, 254
- . 1992a, *Nature*, 359, 619
- . 1992b, *Icarus*, 96, 85
- . 1995, *Icarus*, 117, 287
- . 2012, *Progress of Theoretical Physics Supplement*, 195, 48
- Salo, H. & Karjalainen, R. 2003, *Icarus*, 164, 428
- Salo, H., Karjalainen, R., & French, R. G. 2004, *Icarus*, 170, 70
- Salo, H., Schmidt, J., & Spahn, F. 2001, *Icarus*, 153, 295
- Schmidt, J. & Salo, H. 2003, *Physical Review Letters*, 90, 061102
- Schmidt, J., Salo, H., Spahn, F., & Petzschmann, O. 2001, *Icarus*, 153, 316
- Schmit, U. & Tscharnuter, W. M. 1995, *Icarus*, 115, 304
- Takeda, T. & Ida, S. 2001, *ApJ*, 560, 514
- Tiscareno, M. S., Burns, J. A., Nicholson, P. D., Hedman, M. M., & Porco, C. C. 2007, *Icarus*, 189,

- Toomre, A. 1964, ApJ, 139, 1217
- . 1981, Structure and Evolution of Normal Galaxies (Cambridge: Cambridge Univ. Press), 111
- Toomre, A. & Kalnajs, A. J. 1991, Dynamics of Disc Galaxies, 341
- Wisdom, J. & Tremaine, S. 1988, AJ, 95, 925
- Yasui, Y., Ohtsuki, K., & Daisaka, H. 2012, AJ, 143, 110
- Zebker, H. A., Marouf, E. A., & Tyler, G. L. 1985, Icarus, 64, 531